

Drug	Dose* (mg/kg)	Caution	Comments	Dosing Example
Propofol	2–6	Causes dose-dependent cardiorespiratory depression	Fast onset, short duration, so easy to titrate to effect. Repeat dosing is safe in both cats and dogs	
Alfaxalone	1–3 (D) 2–5 (C)		Fast onset, short duration, easy to titrate to effect. Can be administered IM as a sedative (see sedation chart)	
Etomidate	1–3	Decrease dose by co-inducing with 0.2–0.4 mg/kg of midazolam or diazepam	Can cause myoclonus, paddling, and vocalization at induction but can be prevented with co-induction with a benzodiazepine. Etomidate with propylene glycol is an irritant to veins, causes pain on injection, and perivascular injections causes tissue necrosis/phlebitis. Can cause adrenocortical suppression—use in patients with adrenocortical insufficiency is controversial	Ideal choice for patients with significant cardiovascular disease. Causes minimal to no cardiovascular effects but some respiratory depression with rapid IV administration. Can dilute with fluids running to decrease injection pain and vascular effects
Thiopental	5–15		Will cause tissue necrosis and phlebitis if extravasated; causes respiratory and cardiovascular depression	Currently unavailable in the US
Ketamine + Midazolam or Diazepam	2–5 0.1–0.4	Ketamine used alone causes muscle rigidity and could potentially exacerbate some types of seizures and tachyarrhythmias. These effects are alleviated with the addition of the benzodiazepine. Ketamine is cleared in part unchanged by the kidney in cats so use with caution in cats with renal failure	Most commonly administered as a 1:1 ratio by volume using 100 mg/ml ketamine and diazepam or midazolam (5 mg/ml) with an anticipated total dose of 2 ml/ 20 kg IV in a non-or lightly sedated patient and 1 ml/20 kg IV in a moderately sedated patient	A lightly sedated 20 kg patient would receive 1 ml diazepam + 1 ml ketamine combined in the same syringe for administration
Ketamine + Propofol	2–4 2–4	Same as each drug used individually. May decrease adverse effects of each drug (e.g., less propofol-induced respiratory depression, less ketamine-induced muscle rigidity)	Combine 2–4 mg/kg of each drug (ketamine 100 mg/ml + propofol 20mg/ml) in the same syringe and dose at 2–4 mg/kg of the COMBINATION	For a 20-kg dog, draw up 40 mg (2 mL) propofol and 40 mg (0.4 mL) ketamine. The combination concentration is 80 mg/mL
Tiletamine and Zolazepam	1–4	Without the concurrent use of sedatives , can cause prolonged, “rough,” or dysphoric recovery, especially in dogs. Tiletamine has the same cautions as ketamine and same alleviation of adverse effects because it is combined with a benzodiazepine	Physiologic effects similar to those of ketamine + midazolam or diazepam	These dosages are for a sedated patient. Dosages as high as 6–9 mg/kg may be necessary without sedation

Administer one of the drugs or drug combinations as deemed appropriate for any underlying comorbidity. After sedation is achieved, titrate induction drug “to effect,” anticipating use of the lower end of the dose range for moderately sedated animals or those with moderate systemic disease. A dose lower than that listed may be adequate for heavily sedated, obtunded, severely diseased patients, neonates and “true” geriatrics (i.e., those showing age-related changes). Use caution in patients with inadequate sedation, as this may necessitate a higher dose, which could lead to more physiologic complications.

* Dosages are the same for both dogs (D) and cats (C) unless otherwise indicated.